Anthropogenic particles in the stomach contents and liver of the freshwater fish *Squalius cephalus*

France Collard, Johnny Gasperi, Bernard Gilbert, Gauthier Eppe, Sam Azimi, Vincent Rocher, Bruno Tassin

► To cite this version:

France Collard, Johnny Gasperi, Bernard Gilbert, Gauthier Eppe, Sam Azimi, et al.. Anthropogenic particles in the stomach contents and liver of the freshwater fish *Squalius cephalus*. *Science of the Total Environment*, 2018, 643, pp.1257 - 1264. 10.1016/j.scitotenv.2018.06.313 . hal-01830268

HAL Id: hal-01830268

<https://hal-enpc.archives-ouvertes.fr/hal-01830268>

Submitted on 19 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Anthropogenic particles in the stomach contents and liver of the freshwater fish

2 *Squalius cephalus*

3
4
5
6 France Collard^{1*}, Johnny Gasperi¹, Bernard Gilbert², Gauthier Eppe², Sam Azimi³, Vincent
7 Rocher³, Bruno Tassin¹

8
9
10 ¹Laboratoire Eau Environnement et Systèmes urbains (LEESU), Université Paris-Est Créteil, Ecole
11 des ponts Paristech, AgroParisTech, 61 avenue du Général de Gaulle, 94010 Cedex Créteil, France.

12 ²Inorganic Analytical Chemistry Laboratory, B6c, Department of Chemistry, University of Liege,
13 4000 Liege, Belgium.

14 ³Syndicat Interdépartemental pour l'Assainissement de l'Agglomération Parisienne (SIAAP),
15 Direction du Développement et de la Prospective, 82 Avenue Kléber, 92700 Colombes, France.

16
17 *Corresponding author:

18 Laboratoire Eau Environnement et Systèmes Urbains (LEESU)

19 Université Paris-Est Créteil (UPEC)

20 61, Avenue du Général de Gaulle 94010 Créteil, France

21 francecollard16@gmail.com, gasperi@u-pec.fr

22 phone: + 33145171621

23 Current address: Norwegian Polar Institute (NPI), Fram Center, 9296 Tromsø, Norway

Abstract

Anthropogenic particles (APs) are a very broad category of particles produced directly or indirectly by human activities. Their ingestion by biota is well studied in the marine environment. In contrast, studies on AP ingestion in wild freshwater organisms are scarce despite high contamination levels in some rivers and lakes. In this study, we aimed to evaluate the ingestion of APs and the possible occurrence of APs in the liver and muscle of a freshwater fish, *Squalius cephalus*, from the Parisian conurbation. After isolation, the particles were analyzed using Raman spectroscopy. In sixty stomachs, eighteen APs were found, half of which were plastics and the other half were dyed particles. Twenty-five percent of sampled individuals had ingested at least one AP. The mean length of the APs was 2.41 mm. No significant difference was found between the sites upstream and downstream of Paris. Additionally, 5% of sampled livers contained one or more APs, which were characterized as microplastics (MPs). No APs were found in the muscle tissue. The majority of APs isolated from stomach contents were fibers, which is similar to the findings of a previous river contamination study. This highlights that fish could be more exposed to fibers than previously thought and that more studies on the impacts of fiber ingestion are required. Despite their low occurrence, MPs are reported, for the first time, in the liver of a wild freshwater fish species. While the pathways and impacts are still unknown, MPs also occur in liver of marine mollusks and fish. Physiological *in vitro* studies are needed to better evaluate the impacts of such phenomena.

Keywords

Microplastics, fibers, Seine River, European chub, muscle, liver

Introduction

Anthropogenic particles (APs) are a very broad category of particles produced directly or indirectly by human activities, thus showing an anthropogenic origin. In this study, APs are defined as small pieces – fragments or fibers – with an anthropogenic origin, such as plastic, dyed particles or textile fibers, regardless of their size. All textile fibers have been handled by humans (e.g., dyed) and thus have an anthropogenic origin, regardless of whether their basic composition is natural. All seas and oceans are contaminated by APs, including plastics (Cózar et al., 2014), and so are a large number of organisms (Kühn et al., 2015). Numerous studies, reviews and books (Bergmann et al., 2015) report the extent of this phenomenon, particularly plastic pollution and its impact on marine environments. However, although marine environments have been well studied regarding APs (mainly plastics), knowledge about freshwater contamination by APs and their ecological impacts remains scarce (Dris et al., 2015b). Yet, rivers continue to be the main source of microplastics (MPs) (<5 mm, (Arthur et al., 2009)) to coastal ecosystems (Lima et al. 2014; Lebreton et al. 2017).

Several studies conducted in freshwater (Dris et al., 2015a; Faure et al., 2015), such as in the Seine River (Gasperi et al., 2014), the River Thames (Morritt et al., 2014), the Danube River (Lechner et al., 2014), the Yangtze and Hanjiang Rivers (Wang et al., 2017), the Laurentian Great Lakes (Eriksen et al., 2013) and in several Chilean rivers (Rech et al., 2014), showed that freshwater is contaminated, sometimes highly contaminated, by APs, especially plastics. Among others, a study by Lechner et al. (2014) calculated that, in the Danube River, the global plastic mass for both MPs and macroplastics was higher than the fish larvae mass. In the U.S.A., over 4 million particles are estimated to be released every day from one Californian wastewater treatment plant (Mason et al., 2016).

72

73 High AP concentrations could likely lead to their ingestion by aquatic organisms. Studies focusing
74 on freshwater biota contamination are very scarce. Few studies mention AP or MP ingestion by
75 wild fish species (Faure et al., 2012, 2015; Sanchez et al., 2014; Phillips and Bonner, 2015; Pazos
76 et al., 2017; Silva-Cavalcanti et al., 2017; Vendel et al., 2017) or marine mammal species
77 (Denuncio et al., 2011), and even these have focused only on stomach contents.

78 Aquatic organisms can suffer mechanical damage from ingestion of APs, e.g., obstruction or
79 starvation, as has been shown in marine mammals and birds (Beck and Barros, 1991; Pierce et al.,
80 2004; Jacobsen et al., 2010). Furthermore, aquatic organisms, including fish, can also be impacted
81 by toxicological issues including pollutant transfer (Cedervall et al., 2012; Rochman et al., 2013;
82 Wright and Kelly, 2017) or translocation of MPs into other organs such as the liver (Avio et al.,
83 2015; Collard et al., 2017a).

84

85 In this study, we analyzed the ingestion of APs by the chub *Squalius cephalus* in two highly
86 anthropized rivers, the Marne and the Seine Rivers, which both cross the Greater Paris Megacity.
87 The chub has not been studied so far in the context of AP pollution despite its omnipresence in
88 European rivers and its abundant population (Freyhof, 2014). Due to its ubiquity, the chub could
89 be a bioindicator for AP pollution in rivers. This species has a minor economic interest, but its
90 consumption by Europeans was at its maximum in 2011 (FAO, 2011). Adults are solitary, and
91 juveniles are gregarious, but both are pelagic. Generally, they feed on insects, plants and
92 crustaceans such as crayfish (Balestrieri et al., 2006; Mann, 1976; Michel and Oberdorff, 1995).
93 However, their diet varies with age and season. In the Babuna River (Republic of Macedonia), they
94 mainly feed on algae (Chrysophyceae) and insect larvae (*Ephemeroptera* and *Plecoptera*) in spring,
95 on algae (Chrysophyceae) in summer and autumn and on diatoms and other insect larvae

(Chironomidae) in winter (Nastova-Gjorgjioska et al., 1997). Young individuals eat primarily insects (Mann, 1976), while adults generally feed on plants or crayfish and on other fish, particularly in winter (Michel and Oberdorff, 1995). Moreover, the chub adapts its diet to its habitat (Piria et al., 2005). For example, in Croatia, depending on the river, the diet -dominant components are *Diptera* (insects), cladocerans (Djinova, 1976 cited in Piria et al., 2005), detritus (Adamek and Obrdlik, 1977) or zoobenthos (Losos et al., 1980).

AP contamination of the Greater Paris Megacity rivers, e.g., the Seine and Marne Rivers, was studied during a 19-month period between April 2014 and December 2015, excluding January and August 2015, from upstream to downstream suburbs crossing the very dense part of the urban area (Dris et al., 2018). Because of its sprawling population (12 million inhabitants), Greater Paris exerts a high anthropogenic pressure on rivers and, therefore, is a site of great interest for assessing the ingestion of APs by *Squalius cephalus*. The sampling was done along the continuum of the Seine and Marne Rivers, allowing a comparison between the upstream and the downstream suburbs of Paris. Our objectives were to evaluate the ingestion of APs, including microplastics, by European chubs caught within the Parisian conurbation and to determine whether APs translocate into muscle or liver tissue in this species.

Materials and methods

Sampling

Sixty freshwater fish of the species *Squalius cephalus* (mean length $\pm$ SD: 296 mm $\pm$ 62, mean weight $\pm$ SD: 340 g $\pm$ 239) were electrofished in the Marne and Seine Rivers around Paris (Fig. 1),

France, between 28 August and 2 September 2016. Of the six sampling stations, three were located upstream of Paris (Gournay-sur-Marne, Maisons-Alfort and Villeneuve-Saint-Georges), and three were located downstream of Paris (Triel-sur-Seine, Levallois and Le Pecq). The three target tissues, namely, stomach (n=60), liver (n=60) and muscle (n=22), were immediately removed on board. Stomachs and livers were put in distilled water-diluted formaldehyde solution (5%), while muscles were packaged in aluminum paper and frozen at -20°C in the lab. All tissues were weighed (wet weight) before the isolation process.

Figure 1. Sampling stations in the Parisian conurbation. Circles represent sampling stations and squares represent wastewater treatment plants. Number of stomach content, liver and muscle samples are shown for each station.

Isolation of APs

The entirety of the stomach contents and livers were used, while only a small amount of muscle sample was digested (2-4 g). All tissues were degraded using sodium hypochlorite and then filtered with a 5- μ m filtering membrane according to published methodology (Collard et al., 2015). Nevertheless, methanol (99%), which is typically used for the centrifugation step, was replaced by ethanol (99%), as methanol degrades polyvinylchloride (Collard et al., 2015). Indeed, ethanol is commonly used to store biota samples or microplastics once analyzed (Sanchez et al., 2014; Phillips and Bonner, 2015; Ory et al., 2017). Consequently, in the frame of this study, tissues were put into a 14 g/L NaClO solution overnight. The NaClO solution was filtered with a cellulose acetate filter membrane (5- μ m porosity), which was then rinsed with ethanol. This solution, containing APs, was centrifuged at 5,000 rpm for 10 min. The bottom was then collected and deposited onto a stainless steel plate for Raman spectroscopy analysis.

Raman analysis

All found particles were analyzed using a LabRam 300 spectrometer (Jobin-Yvon) equipped with an Olympus confocal microscope and an Andor BRDD Du401 CCD detector. Depending on the color of the particle, either a Spectraphysics argon ion laser (514.5 nm) or a Torsana diode laser (784.7 nm), with two different objectives (magnification of $\times 50$ or $\times 100$), was used. The maximum beam laser powers induced on the sample were 5 mW (green laser) and 30 mW (red laser), but several neutral density filters were used to decrease the power and avoid degradation of the sample. The integration times ranged from 5 s to 50 s, depending on the sample. Two spectral databases

were used to perform matchings: a commercially available database (Omnics Spectra software, Thermo Fisher Scientific, U.S.A.) and a personal library, which used the Thermo Spectra 2.0 software.

After Raman analyses, APs were isolated in 1 ml of 99% ethanol for further observations and measurements. Only particles whose anthropogenic origin was confirmed by Raman spectroscopy (plastic particles, dyed particles, etc.) were included in the results. Those particles showing an anthropogenic origin have been classified as “APs”. Inside the “AP” category, we defined two subcategories: microplastics and other APs (e.g., dyed particles of an unknown material). In addition, we classified all APs based on their shape: either fragments or fibers.

Preventing contamination and procedural blanks

To minimize contamination, white cotton lab coats and latex gloves were worn throughout the entire isolation process. All work surfaces and dissection materials were rinsed with distilled water, and all glassware was rinsed with distilled water before each use. Stainless steel plates were placed under a metal sifter with 36- μ m mesh to prevent airborne contamination while the plates dried. No particles smaller than 36 μ m were found. The NaClO was filtered with a 5- μ m cellulose acetate membrane before dilution with distilled water.

Three 50-ml volumes of formaldehyde used to store the samples were filtered with the same filter membrane as that used for the isolation process. No APs were found. In addition, guts and livers were kept closed in the formaldehyde but were opened under an airflow cabinet just before the isolation process. Muscles were kept in aluminum, but particles from this material were not found in any samples.

Three procedural blanks were performed, along with runs of several samples, meaning that the entire isolation process was verified. It began in the formaldehyde solution used to store the samples and ended with the observation of stainless steel plates using Raman spectroscopy. This allowed us to evaluate whether solutions, materials or equipment were contaminated by APs in any way. Two particles made of an unknown organic material and two others made of only cellulose were found, meaning that those particles were not APs.

Images and measurements of particles

APs were collected from the Raman plates with 99% ethanol solution and then put into a microtube for further filtration using the same white filter membranes as those used for the isolation process. APs were photographed on both the stainless steel plate and on the filter membrane using a stereomicroscope (Leica MZ12, Leica AG Camera, Germany). They were then measured at their longest dimension using the Histolab software (Histolab Products AB, Sweden).

Statistics

Statistical analyses were performed with the GraphPad Prism software (v5.03, GraphPad software Inc., California, U.S.A.). Pearson's test was applied to evaluate the correlation between fish and AP lengths. Normality of the data was checked using the Kolmogorov-Smirnov test prior to analysis. Fisher's test was applied to compare (1) the number of fish that ingested (or not) at least one AP, MP, fiber or fragment, with each type of particle considered independently; (2) the number of all ingested APs versus MPs; (3) the number of ingested fibers versus fragments; and (4) the concentrations of all APs and MPs in SC, between upstream and downstream of Paris. A Mann-

Whitney U test was applied to compare the lengths of ingested APs between the upstream and downstream suburbs of Paris. All alpha values were set at 0.05. Data are expressed in mean $\pm$ standard deviation (SD).

Results

Stomach contents (SCs)

All isolated particles were analyzed, and 18 APs out of 70 isolated particles were found (Fig. 2). The fifty-two other particles were made of natural organic polymers such as cellulose or could not be identified. Twenty-five percent of fish have ingested at least one AP, and 15% have ingested at least one plastic particle.

Figure 2. Pictures of APs found in stomach contents (A-D) and livers (E, F) of *S. cephalus*. A: blue dyed fiber, B: polypropylene (PP) fiber, C: polyethylene terephthalate fiber, D: PP fiber, E: polystyrene fragment, F: polyethylene fragment. Scale bar: 500 μm .

Table 1. Characteristics of the 18 APs found in stomach contents (SC). *Concerns only fibers.

Station	Composition	Size (mm)	Diameter* (μm)	Shape	Item/g SC
Gournay-sur-Marne	PET	2.96	11	FI	0.38
	Black dyed fiber	0.84	22	FI	2.86
	PAN	7.38	30	FI	
Maisons-Alfort	Blue dyed fragment	2.41	-	FR	0.23

	Blue dyed fiber	0.96	18	FI	
	PET	0.74	12	FI	0.71
Villeneuve-Saint-Georges	PET	2.51	40	FI	2.5
	PP	0.69	59	FI	0.38
	PET	0.39	32	FI	0.67
	Blue dyed fiber	1.08	27	FI	0.38
Triel-sur-Seine	Blue dyed particle	1.43	23	FI	1.7
	Blue dyed fiber	2.21	17	FI	
	PEVA	6.56	-	FR	0.36
	PP	0.44	-	FR	0.67
Levallois	PET	2.36	56	FI	1.0
	Blue dyed fiber	4.44	22	FI	0.59
Le Pecq	Blue dyed fiber	1.21	31	FI	3.33
	Black dyed fiber	4.79	22	FI	2.0

PET=polyethylene terephthalate, PP=polypropylene, PEVA=poly(ethylene-co-vinyl acetate),

PAN=polyacrylonitrile, FI=fiber, FR=fragment.

226

227

Figure 3. Summary of the main characteristics of APs found in stomach contents. a: total AP amount according to shape, b: total AP amount according to composition.

Of the 18 APs, 15 were fibers, and 3 were fragments (Fig. 3a). Fibers longer than 1 mm were dominant (n=10), followed by fibers ranging from 0.5 to 1 mm (n=4) and by fibers smaller than 0.5 mm (n=1). The mean diameter of fibers was $28 \pm 14 \mu\text{m}$. Among the 9 plastic particles, 7 were fibers, and two were fragments. APs measured on average $2.41 \pm 2.09 \text{ mm}$, with a minimum length of $390 \mu\text{m}$ and a maximum length of 7.38 mm. Nine APs were made of plastic polymers and nine were other kinds of APs, i.e. dyed particles (Table 1, Fig. 3b). Four types of plastic polymers were found (Fig. 4): polyethylene terephthalate (PET, n=5), polypropylene (PP, n=2), polyacrylonitrile (PAN, n=1) and poly(ethylene-co-vinyl acetate) (PEVA, n=1). Furthermore, twenty fibers of cellulose were found in SC samples but were not included in APs, because the anthropogenic origin could not be confirmed using Raman spectroscopy. There is no correlation between the length of ingested APs and the length of fish ($R^2=0.0026$).

Figure 4. Raman spectra obtained from isolated APs. PET=polyethylene terephthalate, PP=polypropylene, PEVA=poly(ethylene-co-vinyl acetate), PAN=polyacrylonitrile.

When comparing the areas upstream and downstream of Paris, no significant difference was found regarding the number of individuals having ingested MPs alone, all APs, fibers and fragments; the number of ingested APs and MP; the number of ingested fibers and fragments (contingency tables, $\alpha > 0.05$); and the length of all ingested APs (Mann-Whitney U test, $\alpha = 0.24$). A summary of the results is shown in Table 2. Fish from the upstream stations had 0.32 AP/g of SC, while fish from the downstream stations had 0.40 AP/g of SC, also showing no significant difference (contingency tables, $\alpha > 0.05$).

Table 2. Comparison of different data between the areas upstream and downstream of Paris.
%=percentage.

	Upstream (n=33)	Downstream (n=27)
% of fish with AP	24%	26%
% of fish with MP	18%	11%
Mean AP size	2.00 mm	2.93 mm
Min-max size	0.39-7.38 mm	0.44-6.56 mm
% of fibers	90%	75%
% of fragments	10%	15%
AP/g of SC (wet weight)	0.32	0.40

Livers & Muscles

Four APs, of 13 total extracted particles, were found in three livers out of the sixty analyzed. All were MPs, but three were made of PE, and one was made of polystyrene (PS). Their lengths ranged from 147 μm to 567 μm (Table 3). Given the size of these MP fragments, it is very unlikely that they come from airborne contamination.

Table 3. Characteristics of MPs found in livers.

Station	Polymer	Size (mm)	Shape
Levallois	PE	0.567	FR
	PE	0.161	FR
	PE	0.147	FR

Villeneuve- Saint- Georges	PS	0.270	FR
---	----	-------	----

FR=fragment, PE=polyethylene, PS=polystyrene.

No APs were found in muscle tissue.

Discussion

No difference was found between the areas upstream and downstream of Paris. Regarding shape, fibers were dominant, and fibers longer than 1 mm were found in a higher quantity than the two other size classes. The smallest fiber found in SCs was 390 μm long. Smaller particles are unlikely to be ingested because they would probably be rejected with water through branchial structures (Collard et al., 2017b). Although Dris et al. (2015a) did not use any spectroscopic method for identification, these three results are consistent with those highlighted by the previous study in the Seine River water. Their study found an average of 45 fibers/ m^3 and 0.54 fragments/ m^3 in the water column but found no difference between the downstream and upstream areas, and fibers longer than 1 mm were predominant. European chubs' stomach contents seem to reflect what is found in the river water. Moreover, since it is not an endangered species in European waters, European chub could then be used as a bioindicator for AP and MP pollution. It is already used as a common bioindicator in European freshwaters for other pollutants, such as musk (Hájková et al., 2007), metals (Dragun et al., 2016), dioxins and polychlorobiphenyls (Pacini et al., 2013).

Compared to fibers, only a few anthropogenic fragments were found. Fragments may be indirectly ingested through various pathways: transfer from prey, as shown in experimental studies (Cedervall et al., 2012; Farrell and Nelson, 2013; Setälä et al., 2014); accidental ingestion from the water column or from sediments; or intentional ingestion (Ory et al., 2017). The chub is an omnivorous (Piria et al., 2005) and opportunistic fish (Balestrieri et al., 2006), meaning that everything is potential prey to feed on, including plastic. However, mechanisms of ingestion cannot be elucidated in this study.

All former studies dealing with AP ingestion by freshwater fish focused on MPs. In comparison, the next section will also focus on MP ingestion but not on all APs. Few studies exist on MP ingestion by freshwater fishes, and moreover, their results must be carefully compared. Locations, species, and methodologies, including characterization of the target particles, differ between studies, leading to inconsistent comparisons. Here, we chose to provide data in terms of items/individual, items/g SC and percentage of occurrence. Each of those units has advantages and disadvantages but will at least allow comparisons between previous and future studies. The percentage of MP occurrence is often used to make comparisons more reliable. Regarding MP ingestion by freshwater fish, some of the lowest percentages of MP occurrence are found in fish from Geneva Lake (Faure et al., 2015), from French rivers (Sanchez et al., 2014), from a Brazilian estuary (Vendel et al., 2017), and in European smelts *Osmerus eperlanus* from the Thames estuary (McGoran et al., 2017) (7.5%, 12%, 9%, and 20% of individuals had ingested plastics, respectively). With an MP occurrence rate of 15%, our study has one of the lowest percentages. In addition, some studies have reported high MP occurrence percentages of up to 100% (Pazos et al., 2017). High values were also found in the Thames estuary, where 71% of flounders *Platichthys flesus* have ingested MPs (McGoran et al., 2017). Flounders are a benthic fish, raising questions

about the possible difference between pelagic and benthic feeders. According to a study on European smelts (pelagic) and flounders (benthic), feeding strategies could have an influence on the ingestion of APs (McGoran et al., 2017). The benthic species had a much higher ingestion rate than the pelagic one, but this was not statistically assessed. By contrast, Jabeen et al. (2017) and Murphy et al. (2017) found a significant difference between demersal and pelagic fishes (in freshwaters and marine waters, respectively). The demersal species ingested more APs. However, our study concerned a pelagic species, and we found a relatively low rate of MP occurrence compared to high percentages previously cited. Our percentage of MP occurrence in SC could have been underestimated. Among the dyed particles found in our study, some may have been made of a plastic polymer, which cannot be detected using Raman spectroscopy because of the dye coating. In marine environments, sediments are a sink for microplastics (Woodall et al., 2014). Regarding freshwaters, the literature lacks information about river sediment microplastic contamination, highlighting the need for more studies to assess whether freshwater demersal feeders are more exposed to pelagic ones.

In the SC, a higher proportion of fibers was found compared to fragments (83% vs. 17%), as expected based on other studies (McGoran et al., 2017; Pazos et al., 2017; Silva-Cavalcanti et al., 2017; Vendel et al., 2017). As the digestion protocol may have bleached some particles because of the use of NaClO, some dyed cellulosic fibers may have, in actuality, been dyed textile fibers that were bleached during the isolation process, leading to an underestimation of APs. However, the results concerning the type of particle ingested by chubs are consistent with what is commonly found in the Seine and Marne Rivers: fibers are much more abundant than fragments.

WWTPs could be sources of fibers in urban rivers. While they help to retain a proportion of them, e.g., from 83% to 95% (Dris et al. 2015a), WWTP effluents still constitute as a source of fibers

(Dris et al. 2015a; Leslie et al. 2017). Paris' WWTPs remove long fibers (> 1 mm), but smaller fibers can pass through (Dris et al., 2015a). Atmospheric fallout can also be an additional source of fibers in receiving systems. Indeed, Dris et al. (2016) found that 50% of fibers coming from atmospheric fallout were longer than 1 mm, which is the dominant size class ingested by chubs. In that study, 50% of all fibers were natural, 21% were manufactured by transformation of natural polymers (e.g., rayon), and 29% contain a petrochemical-derived polymer (e.g., mix of cotton and polyamide, or polyurethane). No information regarding the >1 mm size class was given. Even if no direct link can be established between atmospheric fallout and the presence of long fibers in chubs' SC, we speculate that atmospheric inputs in water could be an important source of the longest fibers, which could then be ingested by freshwater biota.

The release of fibers into the environment is already a major concern, since washing machines can release a high amount of fibers per wash (Hartline et al., 2016; Hernandez et al., 2017). A polyester garment can release fibers between 0.033% and 0.039% of its weight (Dubai and Liebezeit, 2013), and six identical fleece garments can release fibers from 0.008% to 0.021% of their total weight (Pirc et al., 2016). Consequently, in urban environments, fibers may be more problematic than fragments, particularly for filtering organisms that have adapted their morphology to retain such particles (Collard et al., 2017b).

An increasing number of studies report AP, mainly MP, ingestion by fish, but the impacts are unknown. Studies which focus on wild-caught species probably have data regarding only a snapshot of the fish's lifetime. After being ingested, MPs are probably excreted (Van Cauwenberghe and Janssen, 2014; Watts et al., 2014; Grigorakis et al., 2017), but they may have a longer retention time than food that is typically (Ward and Kach, 2009; Mazurais et al., 2015). Additionally, no mechanical harm resulting from MPs has been reported in fish, but in birds,

plastics can reduce hunger or cause a reduction in assimilation efficiency (Ryan, 1989; Lenzi et al., 2016). The problem may instead concern pollutants. Microplastic fibers, like other microplastics, adsorb pollutants present in the surrounding water and contain small amounts of additives. They can be transferred to organisms once ingested (Teuten et al., 2009), although this is controversial (Koelmans et al., 2016). Additionally, little is known about adsorption on APs other than those that are plastic. In our study, 9 out of 18 APs were not made of plastic. How do pollutants interact with such particles? Did they also leach from particles to organs? Fibers constitute a major part of AP pollution in freshwater environments, so answering these questions is a priority for AP pollution research.

For the first time, MPs have been found in the livers of wild freshwater fish. Translocation also occurs in Parisian freshwaters, where fragment contamination is lower than fiber contamination (Dris, 2016). Translocation phenomena of such large MP abundances in fish have already occurred in both the laboratory (Avio et al., 2015) and the marine environment (Collard et al., 2017a). Other organisms also exhibit MP translocation (von Moos et al., 2012; Brennecke et al., 2015) but in the laboratory only. In mammals, the uptake of microparticles from the gut and subsequent passage through Peyer's patches in the intestine mucosa has been known for several decades (LeFevre et al., 1978; Jani et al., 1989; Jani et al., 1990). However, fish do not contain Peyer's patches, (Rombout et al., 1993; Cairn and Swan, 2010), and MP translocation pathways through the intestinal barrier are unknown in these organisms. Some hypotheses have been proposed, such as persorption, during which particles pass between cells of the gut epithelium to the circulatory system (Volkheimer, 1975; Wright and Kelly, 2017); endocytosis, which is an internalization of particles in cells of the digestive gland and has been demonstrated in mussels (Von Moos et al., 2012); and, finally, specialized enterocytes called microfold cells may be involved, as suggested

by Browne et al. (2008), again, in mussels. The digestive system in mussels differs significantly from that of fish, but since the literature regarding MP translocation in fish is very scarce, we can only rely on those few studies to piece together an explanation. Considering the large sizes of MPs found in livers in this study, the first hypothesis could be the most relevant. Persorption has been observed for particles of up to 130 μm (Steffens, 1995 cited in Wright and Kelly, 2017). The remaining possibilities seem unlikely considering the sizes of translocated particles: 3 μm (Browne et al., 2008) and smaller than 80 μm (Von Moos et al., 2012). However, this should be confirmed by experimental studies.

No APs were detected in muscle tissue despite the use of the sensitive Raman-based method developed. However, fewer samples, compared to those of SC and liver, were collected, and only several grams of muscle tissue per individual were taken to limit the amount of solution used. Consequently, other studies focusing on contamination of muscle tissue, and of other organs, are needed.

Conclusions

In conclusion, the opportunistic European chub ingested APs in an amount similar to that observed in other freshwater species (McGoran et al., 2017). Our results reflected the main ones highlighted in a study performed in the Seine and Marne Rivers inhabited by chubs. We then suggested that chubs may be used as an indicator species, as they are not endangered but are in fact common throughout Europe. Half of the analyzed APs were found to be made of plastic, highlighting that focusing on plastic particles could underestimate the level of APs ingested, regardless of species. Fibers were the predominant shape found in stomach contents and are ingested with unknown

consequences. Fragments can translocate into other organs by unknown pathways, as shown in other studies. Until now, this was not observed for fibers.

Many questions are yet to be answered. In the field, do MPs translocate into organs other than the liver? What are the pathways of translocation? How do they impact the organism? Does the MP contamination level in the environment influence the translocation process? What is the proportion of ingested plastic that can translocate into the liver? Considering the current global consumption of fish, humans are directly impacted by those issues. Histological analyses could provide answers to these questions. Histological analyses could contribute to give answers to these questions. The next step in AP contamination research could be to focus on *in vitro* studies, which may help to evaluate impacts and determine what type(s) of particles are likely to be translocated and the reasons for such translocation (polymer, size, presence of dyes on surface molecules, etc.).

Acknowledgments

The research leading to these results received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme (FP7/2007-2013) under the REA grant agreement n. PCOFUND-GA-2013-609102, through the PRESTIGE programme coordinated by Campus France (postdoctoral grant). The authors also want to thank the University of Paris-Est Créteil (UPEC) and the SIAAP scientists who helped during the sampling survey.

References

- Adamek, Z., Obrdlik, P., 1977. Food of important cyprinid species in the warmed barb zone of the Oslava river. *Folia Zool.* 26, 171–182.
- Arthur, C., Baker, J., Bamford, H. (eds), 2009. Proceedings of the International Research Workshop on the Occurrence, Effects and Fate of Microplastics Marine Debris.
- Avio, C.G., Gorbi, S., Regoli, F., 2015. Experimental development of a new protocol for extraction and characterization of microplastics in fish tissues: First observations in commercial species from Adriatic Sea. *Mar. Environ. Res.* 111, 18–26.
<https://doi.org/10.1016/j.marenvres.2015.06.014>
- Balestrieri, A., Prigioni, C., Remonti, L., Sgroso, S., Priore, G., 2006. Feeding ecology of *Leuciscus cephalus* and *Rutilus rubilio* in southern Italy. *Ital. J. Zool.* 73, 129–135.
<https://doi.org/10.1080/11250000600679561>
- Barreiros, J.P., Raykov, V.S., 2014. Lethal lesions and amputation caused by plastic debris and fishing gear on the loggerhead turtle *Caretta caretta* (Linnaeus, 1758). Three case reports from Terceira Island, Azores (NE Atlantic). *Mar. Pollut. Bull.* 86, 518–22.
<https://doi.org/10.1016/j.marpolbul.2014.07.020>
- Beck, C.A., Barros, N.B., 1991. The impact of debris on the Florida manatee. *Mar. Pollut. Bull.* 22, 508–510. [https://doi.org/http://dx.doi.org/10.1016/0025-326X\(91\)90406-I](https://doi.org/http://dx.doi.org/10.1016/0025-326X(91)90406-I)
- Bergmann, M., Gutow, L., Klages, M., 2015. Marine anthropogenic litter. Springer.
- Brennecke, D., Ferreira, E.C., Costa, T.M.M., Appel, D., da Gama, B.A.P., Lenz, M., 2015. Ingested microplastics (>100µm) are translocated to organs of the tropical fiddler crab *Uca rapax*. *Mar. Pollut. Bull.* 96, 491–495. <https://doi.org/10.1016/j.marpolbul.2015.05.001>
- Browne, M.A., Dissanayake, A., Galloway, T.S., Lowe, D.M., Thompson, R.C., 2008. Ingested

454 Microscopic Plastic Translocates to the Circulatory System of the Mussel, *Mytilus edulis*
 455 (L.). Environ. Sci. Technol. 42, 5026–5031. <https://doi.org/10.1021/es800249a>

456 Cairn, K., Swan, C., 2010. Fish Physiology: The Multifunctional Gut of Fish, Fish Physiology.
 457 Elsevier Science.

458 Cedervall, T., Hansson, L.-A., Lard, M., Frohm, B., Linse, S., 2012. Food chain transport of
 459 nanoparticles affects behaviour and fat metabolism in fish. PLoS One 7, e32254.
 460 <https://doi.org/10.1371/journal.pone.0032254>

461 Collard, F., Gilbert, B., Compère, P., Eppe, G., Das, K., Jauniaux, T., Parmentier, E., 2017a.
 462 Microplastics in livers of European anchovies (*Engraulis encrasicolus*, L.). Environ. Pollut.
 463 229, 1000–1005. <https://doi.org/10.1016/j.envpol.2017.07.089>

464 Collard, F., Gilbert, B., Eppe, G., Parmentier, E., Das, K., 2015. Detection of Anthropogenic
 465 Particles in Fish Stomachs: An Isolation Method Adapted to Identification by Raman
 466 Spectroscopy. Arch. Environ. Contam. Toxicol. 69, 331–339.
 467 <https://doi.org/10.1007/s00244-015-0221-0>

468 Collard, F., Gilbert, B., Eppe, G., Roos, L., Compère, P., Das, K., Parmentier, E., 2017b.
 469 Morphology of the filtration apparatus of three planktivorous fishes and relation with
 470 ingested anthropogenic particles. Mar. Pollut. Bull. 116, 182–191.
 471 <https://doi.org/10.1016/j.marpolbul.2016.12.067>

472 Cózar, A., Echevarría, F., González-Gordillo, J.I., Irigoien, X., Ubeda, B., Hernández-León, S.,
 473 Palma, A.T., Navarro, S., García-de-Lomas, J., Ruiz, A., Fernández-de-Puelles, M.L.,
 474 Duarte, C.M., 2014. Plastic debris in the open ocean. Proc. Natl. Acad. Sci. U. S. A. 111,
 475 10239–10244. <https://doi.org/10.1073/pnas.1314705111>

476 Denuncio, P., Bastida, R., Dassis, M., Giardino, G., Gerpe, M., Rodríguez, D., 2011. Plastic
 477 ingestion in Franciscana dolphins, *Pontoporia blainvillei* (Gervais and d'Orbigny, 1844),

478 from Argentina. Mar. Pollut. Bull. 62, 1836–1841.

479 <https://doi.org/10.1016/j.marpolbul.2011.05.003>

480 Djinova, L., 1976. Za ishranata za klenot (*Leuciscus cephalus macedonicus* Kar.) i mrenata
 481 (*Barbus cyclolepis strumicae* Kar.) od rekata i od akumulacijata Turija. Folia Balc. 3.

482 Dragun, Z., Tepić, N., Krasnići, N., Teskeredžić, E., 2016. Accumulation of metals relevant for
 483 agricultural contamination in gills of European chub (*Squalius cephalus*). Environ. Sci.
 484 Pollut. Res. 23, 16802–16815. <https://doi.org/10.1007/s11356-016-6830-y>

485 Dris, R., 2016. First assessment of sources and fate of macro- and micro- plastics in urban
 486 hydrosystems: Case of Paris megacity. Université de Paris-Est Créteil.

487 Dris, R., Gasperi, J., Saad, M., Mirande, C., Tassin, B., 2016. Synthetic fibers in atmospheric
 488 fallout: A source of microplastics in the environment? Mar. Pollut. Bull. 104, 290–293.
 489 <https://doi.org/10.1016/j.marpolbul.2016.01.006>

490 Dris, R., Gasperi, J., Rocher, V., Saad, M., Renault, N., Tassin, B., 2015a. Microplastic
 491 contamination in an urban area: A case study in Greater Paris. Environ. Chem. 12, 592–599.
 492 <https://doi.org/10.1071/EN14167>

493 Dris, R., Gasperi, J., Rocher, V., Tassin, B., 2018. Synthetic and non-synthetic anthropogenic
 494 fibers in a river under the impact of Paris Megacity: Sampling methodological aspects and
 495 flux estimations. Sci. Total Environ. 618, 157–164.
 496 <https://doi.org/10.1016/j.scitotenv.2017.11.009>

497 Dris, R., Imhof, H., Sanchez, W., Gasperi, J., Galgani, F., Tassin, B., Laforsch, C., 2015b.
 498 Beyond the ocean: Contamination of freshwater ecosystems with (micro-)plastic particles.
 499 Environ. Chem. 12, 539–550. <https://doi.org/10.1071/EN14172>

500 Dubaish, F., Liebezeit, G., 2013. Suspended microplastics and black carbon particles in the Jade
 501 system, southern North Sea. Water. Air. Soil Pollut. 224.

502 Eriksen, M., Mason, S., Wilson, S., Box, C., Zellers, A., Edwards, W., Farley, H., Amato, S.,
 503 2013. Microplastic pollution in the surface waters of the Laurentian Great Lakes. *Mar.*
 504 *Pollut. Bull.* 77, 177–182. <https://doi.org/10.1016/j.marpolbul.2013.10.007>
 505 FAO, 2011. FAO Capture Production of *Leuciscus Cephalus* [WWW Document]. URL
 506 <http://www.fishbase.se/report/FAO/FAOCatchList.php?scientific=Leuciscus+cephalus>
 507 (accessed 9.15.17).
 508 Farrell, P., Nelson, K., 2013. Trophic level transfer of microplastic: *Mytilus edulis* (L.) to
 509 *Carcinus maenas* (L.). *Environ. Pollut.* 177, 1–3.
 510 <https://doi.org/10.1016/j.envpol.2013.01.046>
 511 Faure, F., Corbaz, M., Baecher, H., de Alencastro, L., 2012. Pollution due to plastics and
 512 microplastics in Lake Geneva and in the Mediterranean Sea. *Arch. Sci.* 65, 157–164.
 513 Faure, F., Demars, C., Wieser, O., Kunz, M., De Alencastro, L.F., 2015. Plastic pollution in
 514 Swiss surface waters: Nature and concentrations, interaction with pollutants. *Environ.*
 515 *Chem.* 12, 582–591. <https://doi.org/10.1071/EN14218>
 516 Freyhof, J., 2014. The IUCN Red List of Threatened Species 2014: *Squalius cephalus* [WWW
 517 Document]. URL <http://www.iucnredlist.org/details/61205/0> (accessed 9.15.17).
 518 Gasperi, J., Dris, R., Bonin, T., Rocher, V., Tassin, B., 2014. Assessment of floating plastic
 519 debris in surface water along the Seine River. *Environ. Pollut.* 195, 163–166.
 520 <https://doi.org/10.1016/j.envpol.2014.09.001>
 521 Grigorakis, S., Mason, S.A., Drouillard, K.G., 2017. Determination of the gut retention of plastic
 522 microbeads and microfibers in goldfish (*Carassius auratus*). *Chemosphere* 169, 233–238.
 523 <https://doi.org/http://dx.doi.org/10.1016/j.chemosphere.2016.11.055>
 524 Hájková, K., Pulkrabová, J., Hajšlová, J., Randák, T., Žlábek, V., 2007. Chub (*Leuciscus*
 525 *cephalus*) as a Bioindicator of Contamination of the Vltava River by Synthetic Musk

526 Fragrances. Arch. Environ. Contam. Toxicol. 53, 390–396. <https://doi.org/10.1007/s00244->
527 006-0190-4

528 Hartline, N.L., Bruce, N.J., Karba, S.N., Ruff, E.O., Sonar, S.U., Holden, P.A., 2016. Microfiber
529 Masses Recovered from Conventional Machine Washing of New or Aged Garments.
530 Environ. Sci. Technol. 50, 11532–11538. <https://doi.org/10.1021/acs.est.6b03045>

531 Hernandez, E., Nowack, B., Mitrano, D.M., 2017. Polyester Textiles as a Source of Microplastics
532 from Households: A Mechanistic Study to Understand Microfiber Release During Washing.
533 Environ. Sci. Technol. 51, 7036–7046. <https://doi.org/10.1021/acs.est.7b01750>

534 Jabeen, K., Su, L., Li, J., Yang, D., Tong, C., Mu, J., Shi, H., 2017. Microplastics and
535 mesoplastics in fish from coastal and fresh waters of China. Environ. Pollut. 221, 141–149.
536 <https://doi.org/http://dx.doi.org/10.1016/j.envpol.2016.11.055>

537 Jacobsen, J.K., Massey, L., Gulland, F., 2010. Fatal ingestion of floating net debris by two sperm
538 whales (*Physeter macrocephalus*). Mar. Pollut. Bull. 60, 765–7.
539 <https://doi.org/10.1016/j.marpolbul.2010.03.008>

540 Jani, P., Halbert, G.W., Langridge, J., Florence, A.T., 1990. Nanoparticle uptake by the rat
541 gastrointestinal mucosa: Quantitation and particle size dependency. J. Pharm. Pharmacol.
542 42, 821–826.

543 Jani, P., Halbert, G.W.W., Langridge, J., Florence, A.T.T., 1989. The Uptake and Translocation
544 of Latex Nanospheres and Microspheres after Oral Administration to Rats. J. Pharm.
545 Pharmacol. 41, 809–812. <https://doi.org/10.1111/j.2042-7158.1989.tb06377.x>

546 Koelmans, A.A., Bakir, A., Burton, G.A., Janssen, C.R., 2016. Microplastic as a Vector for
547 Chemicals in the Aquatic Environment: Critical Review and Model-Supported
548 Reinterpretation of Empirical Studies. Environ. Sci. Technol. 50, 3315–3326.
549 <https://doi.org/10.1021/acs.est.5b06069>

550 Kühn, S., Bravo Rebolledo, E.L., Van Franeker, J.A., 2015. Deleterious Effects of Litter on
 551 Marine Life, in: Bergmann, M., Gutow, L., Klages, M. (Eds.), *Marine Anthropogenic Litter*.
 552 Berlin, p. 447. https://doi.org/http://dx.doi.org/10.1007/978-3-319-16510-3_4
 553 Lebreton, L.C.M., van der Zwet, J., Damsteeg, J.-W., Slat, B., Andrady, A., Reisser, J., 2017.
 554 River plastic emissions to the world's oceans. *Nat. Commun.* 8, 15611.
 555 <https://doi.org/10.1038/ncomms15611>
 556 Lechner, A., Keckeis, H., Lumesberger-Loisl, F., Zens, B., Krusch, R., Tritthart, M., Glas, M.,
 557 Schludermann, E., 2014. The Danube so colourful: A potpourri of plastic litter outnumber
 558 fish larvae in Europe's second largest river. *Environ. Pollut.* 188, 177–181.
 559 <https://doi.org/http://dx.doi.org/10.1016/j.envpol.2014.02.006>
 560 LeFevre, M.E., Vanderhoff, J.W., Laissue, J.A., Joel, D.D., 1978. Accumulation of 2- μ m latex
 561 particles in mouse Peyer's patches during chronic latex feeding. *Experientia* 34, 120–122.
 562 <https://doi.org/10.1007/BF01921939>
 563 Lenzi, J., Burgues, M.F., Carrizo, D., Machín, E., Teixeira-de Mello, F., 2016. Plastic ingestion
 564 by a generalist seabird on the coast of Uruguay. *Mar. Pollut. Bull.* 107, 71–76.
 565 <https://doi.org/10.1016/j.marpolbul.2016.04.016>
 566 Leslie, H.A., Brandsma, S.H., van Velzen, M.J.M., Vethaak, A.D., 2017. Microplastics en route:
 567 Field measurements in the Dutch river delta and Amsterdam canals, wastewater treatment
 568 plants, North Sea sediments and biota. *Environ. Int.* 101, 133–142.
 569 <https://doi.org/10.1016/j.envint.2017.01.018>
 570 Lima, A.R.A., Barletta, M., Costa, M.F., 2015. Seasonal distribution and interactions between
 571 plankton and microplastics in a tropical estuary. *Estuar. Coast. Shelf Sci.* 165, 213–225.
 572 <https://doi.org/10.1016/j.ecss.2015.05.018>
 573 Losos, B., Penaz, M., Kubièkova, J., 1980. Food and growth of fishes of the Jihlava river. *Acta*

574 Sc. Nat. 14, 1–46.

575 Mann, R.H.K., 1976. Observations on the age, growth, reproduction and food of the chub

576 *Squalius cephalus* (L.) in the River Stour, Dorset. J. Fish Biol. 8, 265–288.

577 <https://doi.org/10.1111/j.1095-8649.1976.tb03950.x>

578 Mason, S.A., Garneau, D., Sutton, R., Chu, Y., Ehmann, K., Barnes, J., Fink, P., Papazissimos,

579 D., Rogers, D.L., 2016. Microplastic pollution is widely detected in US municipal

580 wastewater treatment plant effluent. Environ. Pollut. 218, 1045–1054.

581 <https://doi.org/10.1016/j.envpol.2016.08.056>

582 Mazurais, D., Ernande, B., Quazuguel, P., Severe, A., Huelvan, C., Madec, L., Mouchel, O.,

583 Soudant, P., Robbens, J., Huvet, A., Zambonino-Infante, J., 2015. Evaluation of the impact

584 of polyethylene microbeads ingestion in European sea bass (*Dicentrarchus labrax*) larvae.

585 Mar. Environ. Res. 112, 78–85. <https://doi.org/10.1016/j.marenvres.2015.09.009>

586 McGoran, A.R., Clark, P.F., Morritt, D., 2017. Presence of microplastic in the digestive tracts of

587 European flounder, *Platichthys flesus*, and European smelt, *Osmerus eperlanus*, from the

588 River Thames. Environ. Pollut. 220, 744–751. <https://doi.org/10.1016/j.envpol.2016.09.078>

589 Michel, P., Oberdoff, T., 1995. Feeding habits of fourteen European freshwater fish species.

590 Cybium 19, 5–46.

591 Morritt, D., Stefanoudis, P. V, Pearce, D., Crimmen, O.A., Clark, P.F., 2014. Plastic in the

592 Thames: A river runs through it. Mar. Pollut. Bull. 78, 196–200.

593 Murphy, F., Russell, M., Ewins, C., Quinn, B., 2017. The uptake of macroplastic & microplastic

594 by demersal & pelagic fish in the Northeast Atlantic around Scotland. Mar. Pollut. Bull. 122,

595 353–359. <https://doi.org/10.1016/j.marpolbul.2017.06.073>

596 Nastova-Gorgijoska, R., Kostov, V., Georgiev, S., 1997. Nutrition of chub *Leuciscus cephalus*

597 (Linnaeus, 1758) from river Babuna. Ribarstvo 55, 53–65.

598 Ory, N.C., Sobral, P., Ferreira, J.L., Thiel, M., 2017. Amberstripe scad *Decapterus muroadsi*
599 (Carangidae) fish ingest blue microplastics resembling their copepod prey along the coast of
600 Rapa Nui (Easter Island) in the South Pacific subtropical gyre. *Sci. Total Environ.* 586, 430–
601 437. <https://doi.org/10.1016/j.scitotenv.2017.01.175>

602 Pacini, N., Abate, V., Brambilla, G., De Felip, E., De Filippis, S.P., De Luca, S., di Domenico,
603 A., D’Orsi, A., Forte, T., Fulgenzi, A.R., Iacovella, N., Luiselli, L., Miniero, R., Iamiceli,
604 A.L., 2013. Polychlorinated dibenzodioxins, dibenzofurans, and biphenyls in fresh water
605 fish from Campania Region, southern Italy. *Chemosphere* 90, 80–88.
606 <https://doi.org/10.1016/j.chemosphere.2012.07.039>

607 Pazos, R.S., Maiztegui, T., Colautti, D.C., Paracampo, A.H., Gómez, N., 2017. Microplastics in
608 gut contents of coastal freshwater fish from Río de la Plata estuary. *Mar. Pollut. Bull.* 122,
609 85–90. <https://doi.org/10.1016/j.marpolbul.2017.06.007>

610 Phillips, M.B., Bonner, T.H., 2015. Occurrence and amount of microplastic ingested by fishes in
611 watersheds of the Gulf of Mexico. *Mar. Pollut. Bull.*
612 <https://doi.org/10.1016/j.marpolbul.2015.08.041>

613 Pierce, K.E., Harris, R.J., Larned, L.S., Pokras, M.A., 2004. Obstruction and starvation associated
614 with plastic ingestion in a Northern Gannet *Morus bassanus* and a Greater Shearwater
615 *Puffinus gravis*. *Mar. Ornithol.* 32, 187–189.

616 Pirc, U., Vidmar, M., Mozer, A., Kržan, A., 2016. Emissions of microplastic fibers from
617 microfiber fleece during domestic washing. *Environ. Sci. Pollut. Res.* 23, 22206–22211.
618 <https://doi.org/10.1007/s11356-016-7703-0>

619 Piria, M., Treer, T., Aničić, I., Safner, R., Odak, T., 2005. The natural diet of five cyprinid fish
620 species. *Agric. Conspec. Sci.* 70, 21–28.

621 Rech, S., Macaya-Caquilpán, V., Pantoja, J.F., Rivadeneira, M.M., Jofre Madariaga, D., Thiel,

622 M., 2014. Rivers as a source of marine litter - A study from the SE Pacific. Mar. Pollut.
623 Bull. 82, 66–75.

624 Rochman, C.M., Hoh, E., Hentschel, B.T., Kaye, S., 2013. Long-term field measurement of
625 sorption of organic contaminants to five types of plastic pellets: Implications for plastic
626 marine debris. Environ. Sci. Technol. 47, 1646–1654.

627 Rombout, J.H.W.M., Taverne-Thiele, A.J., Villena, M.I., 1993. The gut-associated lymphoid
628 tissue (GALT) of carp (*Cyprinus carpio* L.): An immunocytochemical analysis. Dev. Comp.
629 Immunol. 17, 55–66. [https://doi.org/10.1016/0145-305X\(93\)90015-I](https://doi.org/10.1016/0145-305X(93)90015-I)

630 Ryan, P.G., 1989. The effects of ingested plastic and other marine debris on seabirds, in:
631 Shomura, R.S., Godfrey, M.L. (Eds.), Proceedings of the Second International Conference
632 on Marine Debris. Honolulu, Hawaii, pp. 623–634.

633 Sanchez, W., Bender, C., Porcher, J.-M., 2014. Wild gudgeons (*Gobio gobio*) from French rivers
634 are contaminated by microplastics: preliminary study and first evidence. Environ. Res. 128,
635 98–100. <https://doi.org/10.1016/j.envres.2013.11.004>

636 Setälä, O., Fleming-Lehtinen, V., Lehtiniemi, M., 2014. Ingestion and transfer of microplastics in
637 the planktonic food web. Environ. Pollut. 185, 77–83.
638 <https://doi.org/10.1016/j.envpol.2013.10.013>

639 Silva-Cavalcanti, J.S., Silva, J.D.B., França, E.J. de, Araújo, M.C.B. de, Gusmão, F., 2017.
640 Microplastics ingestion by a common tropical freshwater fishing resource. Environ. Pollut.
641 221, 218–226. <https://doi.org/10.1016/j.envpol.2016.11.068>

642 Steffens, K.-J., 1995. Persorption — Criticism and Agreement as Based upon In Vitro and In
643 Vivo Studies on Mammals, in: Absorption of Orally Administered Enzymes. Springer Berlin
644 Heidelberg, Berlin, Heidelberg, pp. 9–21. https://doi.org/10.1007/978-3-642-79511-4_2

645 Teuten, E.L., Saquing, J.M., Knappe, D.R.U., Barlaz, M.A., Jonsson, S., Bjorn, A., Rowland,

646 S.J., Thompson, R.C., Galloway, T.S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C.,
 647 Viet, P.H., Tana, T.S., Prudente, M., Boonyatumanond, R., Zakaria, M.P., Akkhavong, K.,
 648 Ogata, Y., Hirai, H., Iwasa, S., Mizukawa, K., Hagino, Y., Imamura, A., Saha, M., Takada,
 649 H., 2009. Transport and release of chemicals from plastics to the environment and to
 650 wildlife. *Philos. Trans. R. Soc. B Biol. Sci.* 364, 2027–2045.
 651 <https://doi.org/10.1098/rstb.2008.0284>
 652 Van Cauwenberghe, L., Janssen, C.R., 2014. Microplastics in bivalves cultured for human
 653 consumption. *Environ. Pollut.* 193, 65–70. <https://doi.org/10.1016/j.envpol.2014.06.010>
 654 Vendel, A.L., Bessa, F., Alves, V.E.N., Amorim, A.L.A., Patrício, J., Palma, A.R.T., 2017.
 655 Widespread microplastic ingestion by fish assemblages in tropical estuaries subjected to
 656 anthropogenic pressures. *Mar. Pollut. Bull.* 117, 448–455.
 657 <https://doi.org/10.1016/j.marpolbul.2017.01.081>
 658 Volkheimer, G., 1975. Hematogenous dissemination of ingested polyvinyl chloride particles.
 659 *Ann. N. Y. Acad. Sci.* 246, 164–71.
 660 Von Moos, N., Burkhardt-Holm, P., Köhler, A., 2012. Uptake and effects of microplastics on
 661 cells and tissue of the blue mussel *Mytilus edulis* L. after an experimental exposure. *Environ.*
 662 *Sci. Technol.* 46, 11327–11335.
 663 Wang, W., Ndungu, A.W., Li, Z., Wang, J., 2017. Microplastics pollution in inland freshwaters
 664 of China: A case study in urban surface waters of Wuhan, China. *Sci. Total Environ.* 575,
 665 1369–1374. <https://doi.org/10.1016/j.scitotenv.2016.09.213>
 666 Ward, J.E., Kach, D.J., 2009. Marine aggregates facilitate ingestion of nanoparticles by
 667 suspension-feeding bivalves. *Mar. Environ. Res.* 68, 137–142.
 668 <https://doi.org/10.1016/j.marenvres.2009.05.002>
 669 Watts, A.J.R., Lewis, C., Goodhead, R.M., Beckett, S.J., Moger, J., Tyler, C.R., Galloway, T.S.,

670 2014. Uptake and retention of microplastics by the shore crab *Carcinus maenas*. Environ.
671 Sci. Technol. 48, 8823–8830.

672 Woodall, L.C., Sanchez-Vidal, A., Canals, M., Paterson, G.L.J., Coppock, R., Sleight, V.,
673 Calafat, A., Rogers, A.D., Narayanaswamy, B.E., Thompson, R.C., 2014. The deep sea is a
674 major sink for microplastic debris. R. Soc. Open Sci. 1, 140317–140317.
675 <https://doi.org/10.1098/rsos.140317>

676 Wright, S.L., Kelly, F.J., 2017. Plastic and Human Health: A Micro Issue? Environ. Sci. Technol.
677 51, 6634–6647. <https://doi.org/10.1021/acs.est.7b00423>
678
679